

## SCD's Discharge Instructions

Please use the compression stockings that were sent home with you to help prevent blood clots after surgery. Plan on using them for 3-5 days post surgery. Use them at bedtime and anytime during the day when you are relaxing (sitting, you should have your legs elevated). Walking is still the best prevention, so only use the stockings at times of rest.

Once you have returned to your pre-surgery activity level, you can return the device in the prepaid envelope, which is inside of your box. Also the lid of the box has the instructions on how to return it as well. If you have questions on how to operate the system please call 989-213-6733.

**COMPRESSION SOLUTIONS**

Check out our How-To Video at [www.CompressionSolutions.us](http://www.CompressionSolutions.us)

Please call 800-994-0464 with any questions.

### Quick Start Guide

- Apply the DVT calf wrap(s) to your leg(s) as shown.
 

→ Air Tube (Back of Calf)
- Plug the hose(s) into the ports by depressing the metal thumb tabs on the port at the bottom of the machine. Plug the AC adapter into the BATTERY CHARGER INPUT.
 

Power Adapter

Leg One

Leg Two

If only using one calf wrap, plug hose into Leg 1.
- Turn the machine on by pressing POWER button until the lights are illuminated.
 

Press Power
- Press and release the MODE button to illuminate Leg 1 and/or Leg 2 and/or AUX, matching the ports where the wraps are plugged.
 

Select Mode (Leg 1, Leg 2 or AUX)

WARNING: Remove leg wraps before walking.